

# NESS & BEAULY FISHERIES TRUST

---

## Trout (*Salmo trutta*) as predators of juvenile Atlantic salmon (*Salmo salar*): a literature review.



A document prepared by the Ness & Beaully Fisheries Trust

*Version 1.1 (24/3/2017)*



## DOCUMENT CONTROL

Created by	Date	Signed
Nick Barker (Senior Fisheries Biologist, NBFT)	24/3/17	
Reviewed By	Date	Signed
Chris Conroy (Director, NBFT)	24/3/17	
Approved by	Date	Signed
Chris Conroy (Director, NBFT)	24/3/17	
File Reference	Revision	Notes
C:\Users\NBFT-1\Documents\Research Journals	Version 1.1	

### Ness & Beaully Fisheries Trust

Corff House  
Beaully  
Inverness-shire  
IV4 7BE

Telephone: 01463783505

Email: [nessandbeaully@gmail.com](mailto:nessandbeaully@gmail.com)

Directors: Neil Cameron (Chairman), Michael Martin, Jock Miller, Graham Mackenzie, James Braithwaite, Frank Spencer-Nairn, Murray Stark.

Registered Office: Harper Macleod, Alder House, Cradlehall Business Park, Inverness, IV1 1YN

Company Number: SC294401

Charity Number: SC037

## Table of Contents

<b>1</b>	<b>Introduction .....</b>	<b>1</b>
<b>2</b>	<b>The Sea Trout .....</b>	<b>1</b>
<b>3</b>	<b>The Diet of Brown Trout .....</b>	<b>2</b>
<b>4</b>	<b>The Diet of Sea Trout .....</b>	<b>3</b>
<b>5</b>	<b>Discussion and Conclusions .....</b>	<b>4</b>

## 1 INTRODUCTION

In 2016, the Lower Beaully Fishing Syndicate enjoyed their highest catch of sea trout (*Salmo trutta*) for many years. Indeed, catches of sea trout outnumbered salmon landings by a ratio of 1.9:1. To the author's knowledge, this is the first time this has happened in living memory. Whilst this news has been welcomed by many, there have been some river users who have aired concerns that this apparent sudden influx of sea trout may have negative impacts on the established population of Atlantic salmon (*Salmo salar L.*) through active predation on juveniles. To this end, The Ness and Beaully Fisheries Trust (NBFT) agreed to review the current literature on the aforementioned subject and produce a report documenting their findings.

## 2 THE SEA TROUT

Sea trout are the anadromous (migratory) form of the brown trout. Anadromous trout are not as extensively distributed as inland populations of brown trout but are found in Iceland, Scandinavia as well as the Baltic and North Sea and as far south in the Atlantic as the Bay of Biscay (Frost & Brown 1967). Early life history traits of sea trout follow that of the brown trout and salmon; adults spawn in the late autumn/winter in small and large waterbodies with the resultant ova developing over winter and hatch in the subsequent spring. Following a period of endogenous (yolk sack consumption period) feeding during the larval (alevin) stage, the resultant fry begin to actively feed on aquatic invertebrates. Following a period of two-three years, some trout may become quite sedentary in rivers but can often appear to move about river catchments freely (Gowan *et al* 1994).

Brown trout occurring in rivers with unhindered access to the sea can form anadromous populations and will leave their natal streams in the spring for coastal waters in a process termed 'smoltification'. It is at this point that brown trout become sea trout. According to Nall (1929), sea trout of Beaully origin do not venture further than the Inner Moray Firth. The migration to sea may be for one summer only before returning as 'finnock'. These small sea trout (typically weighing 0.5 to 0.75 lb) will return to their natal stream but may not spawn during this time unlike larger (multi-sea winter) sea trout who will spawn with their counterparts close to, or in the waterbody from which they were spawned (Fahy 1978). Some sea trout perish post-spawning although some individuals of Beaully origin may spawn up to four times (Nall 1929).

### 3 THE DIET OF BROWN TROUT

The brown trout is widely accepted as an opportunistic feeder with the diet varying with fish size, age, habitat and season (Bridcut & Giller 1995). The diet of stream dwelling brown trout is well documented (Hernandez & Cobo 2015, Jonsson & Gravem 1985). As young parr and fry, insect larvae of chironomids can be the main food item but surface arthropods such as flying insects may also be taken. With increasing size, trout will gradually move on to larger food items such as the larvae and adult forms of the groups Ephemeroptera (mayfly) and Trichoptera (caddis). Grey (2001) documented aspects of dietary specialisation rather than opportunism in stream dwelling trout with some individuals targeting one food item (e.g. chironomid larvae) whilst seemingly ignoring other food items.

Loch dwelling trout may even turn wholly piscivorous and will actively predate on other trout and species such as char (*Salvelinus alpinus*) and three-spined stickleback (*Gasterosteus aculeatus*) (Grey 2001). Piscivory in river dwelling trout in the UK is less well documented although this has been recorded in Norway (Forseth & Jonsson 1994), Ireland (Keeley & Grant 2001) and New Zealand (McIntosh 2011). However, anecdotal evidence from Mckelvey 2017 (personal communication, February 2017) and Walker 2017 (personal communication, March 2017) suggests that brown dwelling in and around hydro-electric storage reservoirs can actively predate on Atlantic salmon smolts during their outward migration. These areas act as 'pinch points' where smolts congregate at entrances to Borlund fish lifts or turbine intakes. Furthermore, in 'truck and transport' operations conducted on the River Conon, brown trout have exhibited habitual behaviour; congregating in areas where salmon smolts will be transported to and actively predated on disorientated individuals (Mckelvey 2017, personal communication February 2017) upon release.

These communications are in line with work published by Piggins (1962) who concluded that brown trout of 230-320mm in length were "serious" predators of salmon smolts in Ireland whilst Mills (1964) noted brown trout of the same length consumed salmon fry between April and November in Ross-shire. Predation on salmon fry by brown trout has also been reported by Skaala *et al* (2014) in a Norwegian river system. More recent research conducted on the River Conon showed in-river mortality of salmon smolts to be in the order of 20% and this was partly attributed to predation by brown trout (Mckelvey 2017, personal communication February 2017).

## 4 THE DIET OF SEA TROUT

Feeding habits of sea trout in the marine environment has been extensively studied (Pemberton 1976, Fahy 1983, Grovnik & Klemetsen 1987, Lyse *et al* 1998, Knutsen *et al* 2001) and more generally by Nall (1930) and Went (1962) as reviewed by Elliot *et al* (1992). In a study from a Scottish Sea Loch, it was found that sea trout took about 50 prey species, according to the contents of 1,277 stomachs examined over one year (Pemberton 1976). In larger individuals of >21cm, clupeid fishes such as herring (*Clupea harengus*), sprat (*Sprattus sprattus*) and sand eel (*Hyperoplus lanceolatus*) were the dominant food whilst smaller fish of <21cm were seen to feed on small crustaceans and insects. These findings were largely confirmed by Fahy (1985) although this study found sea trout to be feeding on 60 different prey items including sprats, sand eels, stickleback and sand smelts (*Atherina presbyter*).

Prior to a study looking at the impact of marine fish predation on salmon smolts in the Tana Estuary, Norway; sea trout were assumed to be (along with other marine species) major predators of salmon smolts in the River Estuary (Svenning *et al* 2005). However, their findings confirmed this was not the case as salmonid remains (were absent from the stomach contents of 56 adult sea trout caught over the space of around two months. These results are in contrast to the findings of Vollset *et al* (2016) who found salmon smolts in low numbers in the stomachs of adult sea trout. In the opinion of Mahlum 2017 (personal communication, February 2017) it is unlikely sea trout are having a large impact on the population of salmon and that the predation on salmon smolts was much more pronounced in species such as cod (*Gadhus morhua*) and pollack (*Pollachius virens*).

There is very little information on the food of adult sea trout after they have migrated into fresh water (Elliot *et al* 1992). Indeed, some sources would suggest that sea trout may not feed at all in fresh water (Wild Trout Trust 2017). Early information summarised by Harris (1971) concluded that Nall (1926, 1930) was the only worker to examine the stomachs of sea trout caught above the tidal limit. Nall's analysis of 150 sea trout ranging from 0.5lb to 3.5lb in weight indicated a diet of terrestrial insects, aquatic insects and crustaceans but no fish. Harris (1971) examined the stomachs of 150 sea trout caught in a Welsh river between 1968 and 1969. These fish were seen to have been in freshwater for up to three weeks; 2% contained any appreciable amount of food with 75% (112 stomachs) were recorded as empty. Of those 2%, stomach contents were a mixture of terrestrial and freshwater invertebrates together with a small number of salmonid eggs and parr.

Elliot (1996) conducted a much larger study of 467 sea trout stomachs (246 males, 221 females) from six NW England rivers ranging from 0.35lb and 2lb in weight. Fish were taken at a range of sampling reaches from 2km to 25km from the tidal limit. 64% of the samples exhibited empty stomachs whilst those captured at a range of 2km to 12km from the tide had remains of marine species in their stomachs such as clupeid fishes, brown shrimp (*Crangon vulgaris*) and shore crab (*Carcinus maenus*). In the rivers furthest from the tidal limit, there was a marked change in diet from marine to freshwater and terrestrial food. No salmonid fish were identified in any of the stomachs despite each river containing a viable population of both salmon and trout. To the author's knowledge, this is the only peer reviewed study on the stomach contents of sea trout caught in fresh water.

## 5 DISCUSSION AND CONCLUSIONS

This document has investigated the currently available peer reviewed literature on the subject of sea trout as predators of Atlantic salmon. Given the anadromous nature of trout, it has also included references to the non-anadromous trout as predators of salmon.

Whilst evidence would suggest that sea trout will predate on juvenile salmon in both the freshwater and marine environment; those authors who have specifically targeted research in this area have concluded that effects on established populations of salmon would be negligible (Mahlum 2017) (personal communication, February 2017) and largely agree with Nall (1926) who concluded that *"There is little doubt that after leaving salt water sea trout feed more intermittently and with a less robust appetite than brown trout"*.

With a rise in numbers of sea trout within the Beaully catchment, there is little doubt that densities of juvenile trout will increase and this is already being seen, particularly in the main tributary burns of the Lower River Beaully; namely the Belladrum Burn, Bruiach Burn and Culburnie Burn. Young salmon and trout forage for the same food items and will utilise similar habitats during the fry stage. Therein lies the question of whether an increase in juvenile trout will impact on juvenile salmon through increased competition in the nursery areas.

This has been addressed and debated for many years (Heggberget *et al* 1988, Kalleberg 1958, Kennedy & Strange 1986). Most recently this was addressed by Milner *et al* (2006) who stated *"... in spite of unambiguous evidence of interspecific competition, there is as yet no clear demonstration of negative*

*interactions between salmon and sea trout populations at fishery or catchment management scales... the possibility of such effects cannot be eliminated... because the required studies in to combined species stock-recruitment relationships have never been carried out and this represents a continuing research need".*

## REFERENCES

- Bridcutt E.E., Giller P.S., 1995. *Diet variability and foraging strategies in brown trout (Salmo trutta): an analysis from subpopulations to individuals.* Canadian Journal of Fisheries and Aquatic Sciences. 52(12) pp 2543-2552.
- Elliot J.M., Crisp D.T., Mann R.H.K., Pettman I., Pickering A.D., Pottinger T.G., Winfield I.J., 1992. *Sea trout literature review and bibliography.* National Rivers Authority: Fisheries Technical Report No.3, pp 1-141.
- Fahy E., 1983. *Food and gut parasite burden of migrant trout (Salmo trutta L.) in the sea.* The Irish Naturalist's Journal. Vol 21(1) pp 11-18
- Fahy E. 1978. *Variation in some biological characteristics of British Sea trout (Salmo trutta L.).* Journal of Fish Biology. 13(1) pp 123-138.
- Forseth T., Jonsson B., 1994. *The growth and food ration of piscivorous brown trout (Salmo trutta).* Functional Ecology. Vol. 8(2) pp 171-177.
- Frost W.E., Brown M.E. 1967. *The Trout.* Collins, London. 286pp.
- Gowan C., Young M.K., Fausch K.D., Riley S.C. 1994. *Restricted movements in resident stream salmonids: A paradigm lost?* Canadian Journal of Fisheries and Aquatic Sciences. 51(11) pp 2626-2637.
- Grovnik S., Klemetsen A., 1987. *Marine food and diet overlap of co-occurring Arctic char (Salvelinus Alpinus L.) Brown trout (Salmo trutta L.) and Atlantic salmon (Salmo salar L.) off Senja, N. Norway.* Polar Biology. 7(173) pp 173-177.
- Harris G.S., 1971. *The freshwater feeding of adult sea trout in the Afon Dyfi (mid-Wales).* Fisheries Management. Vol.2 pp 20-23.
- Heggberget T.G., Haukebo T., Mork J., Stahl G., 1988. *Temporal and spatial segregation of spawning in sympatric populations of Atlantic salmon (Salmo salar L.) and brown trout (Salmo trutta L.).* Journal of Fish Biology. 33 pp 347-356.
- Jonsson B., Gravem F.R., 1985. *Use of space and food by resident migrant brown trout, Salmo trutta.* Environmental Biology of Fishes. Vol. 14(4) pp 281-293.
- Grey J., 2001. *Ontogeny and dietary specialisation in brown trout (Salmo trutta L.) from Loch Ness, Scotland, examined using stable isotopes of carbon and nitrogen.* Ecology of Freshwater Fish. 10 pp 168-176.
- Kalleberg H., 1958. *Observations in a stream tank of territoriality and competition in juvenile salmon and trout.* Institute of Freshwater Research Drottingholm. Report no. 39, pp 55-98.
- Keeley E.R., Grant J.W.A., 2001. *Prey size of salmonid fishes in streams, lakes and oceans.* Canadian Journal of Fisheries and Aquatic Sciences. 58(6) pp 1122-1132.

Kennedy G.J.A., Strange C.D., 1986. *The effects of intra and inter specific competition on the distribution of stocked juvenile Atlantic salmon, Salmo salar L., in relation to depth and gradient in an upland trout, Salmo trutta L., stream.* Journal of Fish Biology. 29 pp 199-214.

Knutsen J.A., Knutsen H., Olsen E.M., Jonsson B., 2004. *Marine Feeding of anadromous Salmo trutta during winter.* Journal of Fish Biology. 64 pp 89-99.

Lyse A.A, Stefansson S.O, Ferno A. 1998. *Behaviour and diet of Sea trout post smolts in a Norwegian Fjord System.* Journal of Fish Biology. Vol 52(5) pp 923-936

McIntosh A.R., 2000. *Habitat and size related variations in exotic trout impacts on native galaxid fishes in New Zealand streams.* Canadian Journal of Fisheries and Aquatic Sciences. 57(10), pp 2140-2151

Mills D.H., 1964. *The ecology of the young stages of the Atlantic salmon in the River Bran, Ross-shire.* Freshwat. Salm.Fish.Res. 32, 58pp

Milner N.J., Karlsson L., Degerman E., Johlander A., Maclean J.C., Hansen L-P., 2006. *Sea Trout (Salmo trutta L.) in European Salmon (Salmo salar L.) Rivers.* In: *Sea Trout, Biology, Conservation & Management.* Proceedings of the First International Sea Trout Symposium, Cardiff, July 2004. Blackwell Publishing Professional, Iowa, USA.

Nall H.G 1929. *Sea trout from the Beaully Firth and from the tidal waters of the Beaully and Ness Rivers.* Fisheries, Scotland, Salmon Fish. No. 111 pp 3-41.

Nall G.H., 1930. *The life of sea trout especially in Scottish Waters; with chapters on the reading and measuring of scales.* Seeley, Servive and Co Ltd.

Pemberton R., 1976. *Sea trout in North Argyll Sea Lochs: II. Diet.* Journal of Fish Biology. Vol 9 pp 195-208.

Piggins D.J., 1962. *Thyroid feeding of salmon parr.* Nature. Vol 195(4845), pp 1017-1018

Sanchez Hernandez J., Cobo F., 2015. *Adaptive flexibility in the feeding behaviour of brown trout: optimal prey size.* Zoological Studies pp54:26

Skalla O., Glover K.A., Barlaup B.T., Borgstrom R. 2014. *Microsatellite DNA used for parentage identification of partly digested Atlantic salmon (Salmo salar) juveniles through non-destructive diet sampling in salmonids.* Marine Biology Research. Vol. 10(3) pp 323-328.

Went A.E.J 1962. *Irish sea trout: A review of investigations to date.* Royal Dublin Society.

Wild Trout Trust 2017 [online]. *Sea Trout.* Available at: <http://www.wildtrout.org/content/sea-trout> [Accessed 20/3/17].